



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

II. Solution by J. C. NAGLE, M. A., M. C. E., Professor of Civil Engineering and Physics, Agricultural and Mechanical College, College Station, Texas.

If the point of projection be origin and the path be regarded as a parabola its equation will be

$$y = x \tan \alpha - \frac{gx^2}{2v^2 \cos^2 \alpha} \dots (1),$$

where α = angle of elevation.

At the point where the shell is to strike the plane, $y = -3750$ feet and $x = 6495$ feet. Inserting these values in equation (1) there results

$$-3750 \times 400 \cos^2 \alpha = 6495 \times 200 \times \sin \alpha \cos \alpha - (1299)^2.$$

For $\cos^2 \alpha$ write $1 + \cos 2\alpha$, and for $2 \sin \alpha \cos \alpha$ write $\sin 2\alpha$, and as $\sin 2\alpha = \sqrt{1 - \cos^2 2\alpha}$, we get

$$\sqrt{1 - \cos^2 2\alpha} = .722 - .577 \cos 2\alpha \dots (2).$$

Solving (2) we get $\cos 2\alpha = 314 \pm .676$.

$\therefore 2\alpha = 8^\circ 48'$ or $111^\circ 12'$ and $\alpha = 4^\circ 24'$ or $55^\circ 36'$, which values satisfy the condition

$$\alpha'' - \frac{1}{2}(\frac{1}{2}\pi - 30) = \frac{1}{2}(\frac{1}{2}\pi - 30) - \alpha'.$$

See Tait and Steele's *Dynamics of a Particle*, page 90.

Also solved by P. H. PHILBRICK.

78. Proposed by ALOIS F. KOVARIK, Professor of Mathematics, Decorah Institute, Decorah, Iowa.

A cone and a cylinder having equal heights and equal circular bases are filled with water; if they have equal holes in the bases, respectively, how many times as long will it take the cylinder to empty as the cone?

I. Solution by P. H. PHILBRICK, C. E., Lake Charles, La.

Let r = radius of base, h = altitude, k = area of orifice, and x = height of the water at the time t .

I. For the cylinder, the discharge in time dt is, $k\sqrt{2gx}dt$; and since in the same time the surface of the water descends a distance dx , the quantity in the vessel is lessened $\pi r^2 dx$.

$\therefore k\sqrt{2gx}dt = \pi r^2 dx$, and

$$t = -\frac{\pi r^2}{k\sqrt{2g}} \int \frac{dx}{\sqrt{x}} = -\frac{2\pi r^2}{k\sqrt{2g}} x^{\frac{1}{2}} + c = \frac{2\pi r^2}{k\sqrt{2g}} (h^{\frac{1}{2}} - x^{\frac{1}{2}}).$$

II. For the cone, we have $y = (r/h)(h-x)$, and the area of section

$$= \pi y^2 = \frac{\pi r^2}{h^2} (h-x)^2.$$

$$\begin{aligned}\therefore k \sqrt{2gx} dt &= \frac{\pi r^2}{h^2} (h-x)^2 dx, \text{ and } t = -\frac{\pi r^2}{h^2 k \sqrt{2g}} \int \frac{(h-x)^2}{x^{\frac{3}{2}}} dx \\ &= -\frac{\pi r^2}{h^2 k \sqrt{2g}} \int \frac{(h^2 - 2hx + x^2) dx}{x^{\frac{3}{2}}} \\ &= -\frac{\pi r^2}{h^2 k \sqrt{2g}} \cdot (2h^2 x^{\frac{1}{2}} - \frac{4h}{3} x^{\frac{3}{2}} + \frac{2}{5} x^{\frac{5}{2}}) + c.\end{aligned}$$

But $t=0$ for $x=h$. $\therefore c = \frac{\pi r^2}{h^2 k \sqrt{2g}} \cdot \frac{16}{15} h^{\frac{5}{2}}.$

Hence $t = -\frac{\pi r^2}{h^2 k \sqrt{2g}} (2h^2 x^{\frac{1}{2}} - \frac{4h}{3} x^{\frac{3}{2}} + \frac{2}{5} x^{\frac{5}{2}}) + \frac{\pi r^2}{k \sqrt{2g}} \cdot \frac{16}{15} h^{\frac{5}{2}}.$

For $x=0$, the time of emptying vessel $= t_2 = \frac{\pi r^2}{k \sqrt{2g}} \cdot \frac{16}{15} h^{\frac{5}{2}}.$

This is $\frac{8}{15}$ of the time of emptying the cylinder.

II. Solution by C. HORNING, A. M., Professor of Mathematics, Heidelberg University, Tiffin, Ohio.

Let A = the area of the descending surface, O = the area of the orifice, and x = the depth of the water at the end of any time t .

The quantity of water discharged through the orifice in the infinitely small time dt is $O \cdot dt(2gx)^{\frac{1}{2}}$, the velocity of discharge being $(2gx)^{\frac{1}{2}}$; but in the same time the surface has descended through the distance dx and the quantity discharged is $A dx$.

$\therefore O \cdot dt(2gx)^{\frac{1}{2}} = A dx$, or $dt = \frac{A dx}{O \cdot (2gx)^{\frac{1}{2}}}$, as general formula for any shape of vessel.

Now for the cylinder $A = \pi r^2$ and therefore $dt = -\frac{\pi r^2 dx}{O(2gx)^{\frac{1}{2}}}$ and

$$t = -\frac{\pi r^2}{O(2g)^{\frac{1}{2}}} \int_h^0 \frac{dx}{x^{\frac{1}{2}}} = \frac{2\pi r^2 h^{\frac{1}{2}}}{O(2g)^{\frac{1}{2}}} = \frac{\pi r^2 (2h)^{\frac{1}{2}}}{O \cdot g^{\frac{1}{2}}};$$

and for the cone $A = \frac{\pi r^2 (h-x)^2}{h^2}$ and

$$\therefore dt = -\frac{\pi r^2 (h-x)^2 dx}{O h^2 (2g)^{\frac{1}{2}}} \text{ and } t = -\frac{\pi r^2}{O h^2 (2g)^{\frac{1}{2}}} \int_h^0 \frac{(h-x)^2 dx}{x^{\frac{1}{2}}}$$

$$= \frac{\pi r^2}{O h^2 (2g)^{\frac{1}{2}}} \cdot \frac{16 h^{\frac{5}{2}}}{15} = \frac{8}{15} \cdot \frac{\pi r^2 (2h)^{\frac{1}{2}}}{O \cdot g^{\frac{1}{2}}}.$$

Comparing the two results we see that the cone empties in $\frac{8}{15}$ of the time it takes the cylinder, or the cylinder takes $1\frac{1}{2}$ as long as the cone to empty. The minus sign is prefixed because x decreases as t increases.

Also solved by G. B. M. ZERR, ELMER SCHUYLER, J. SCHEFFER, and J. C. NAGLE.

NOTE. In reference to problem 63, Dr. Arnold Emch says: "I had the problem solved by my class in graphic statics by a purely graphical method and the following values (approximations) were obtained: $\angle ABE=46^\circ$, $\angle BAD=56^\circ$, tension in $BE=56.8$, tension in $AD=71$. This shows that the solution in the MONTHLY is correct."

DIOPHANTINE ANALYSIS.

71. Proposed by A. H. BELL, Hillsboro, Ill.

Find five numbers such that the product of any two plus 1 will equal a square.

III. Solution by M. A. GRUBER, A. M., War Department, Washington, D. C.

By using $(s-1)^2$, the denominator of Euler's fifth number, where $s=4n(n-1)(n+1)[4n(2n-1)(2n+1)]$, I have found five numbers in terms of n : $x=n-1$, $y=n+1$, $z=4n$, $w=4n(2n-1)(2n+1)$, and

$$v = \frac{4n(2n-1)(2n+1)[2n(2n-1)-1][2n(2n+1)-1](8n^2-1)}{\{4n(n-1)(n+1)[4n(2n-1)(2n+1)]-1\}^2}.$$

The numerator of v is four times the product of the roots of the six squares $xy+1$, $xz+1$, $yz+1$, $xw+1$, $yw+1$, and $zw+1$.

The denominator of v is the square of $(xyzw-1)$.

Take $n=1, 2, 3, 4, 5, 6$, etc. We then obtain the following sets of five numbers:

$$\begin{array}{lllll} 0, & 2, & 4, & 12, & 420; \\ 1, & 3, & 8, & 120, & \frac{777480}{(2879)^2}; \\ 2, & 4, & 12, & 420, & \frac{35455980}{(40319)^2}; \\ 3, & 5, & 16, & 1008, & \frac{499902480}{(241919)^2}; \\ 4, & 6, & 20, & 1980, & \frac{3822388020}{(950399)^2}; \\ 5, & 7, & 24, & 3432, & \frac{20000100120}{(2882879)^2}; \text{ etc., etc., etc.} \end{array}$$

We also find